

Acceleration at $z > 1$?

Luca Amendola

*INAF - Osservatorio Astronomico di Roma, Via Frascati 33, 00040 Monte Porzio Catone (Roma) Italy,
amendola@mporzio.astro.it*

Accepted, Received

ABSTRACT

Most models of dark energy predict the beginning of the accelerated epoch at $z \leq 1$. However, there are no observational or theoretical evidences in favor of such a recent start of the cosmic acceleration. In fact, a model of dark energy coupled to dark matter is explicitly constructed that *a*) is accelerated even at high z ; *b*) allows structure formation during acceleration; and *c*) is consistent with the type Ia supernovae Hubble diagram, including the farthest known supernova SN1997ff at $z \approx 1.7$. It is shown that the accelerated epoch in this model could have started as early as $z \approx 5$.

Key words:

1 INTRODUCTION

Cosmic acceleration is one of the most exciting discoveries of the recent years. The combined data from cosmic microwave background (CMB, see Netterfield et al. 2001, Lee et al. 2001, Halverson et al. 2001), cluster masses and abundance (see e.g. Bahcall et al. 2002) and supernovae Type Ia (SNIa) Hubble diagrams (Riess et al. 1998, Perlmutter et al. 1999) indicate that the universe is currently dominated by a very weakly clustered component that is able to accelerate the expansion. This component, denoted dark energy or quintessence, is supposed to fill roughly 70% of the cosmic medium.

However, the nature of the dark energy is still enigmatic (Wetterich 1988; Ratra & Peebles 1988; Frieman et al. 1995; Caldwell et al. 1998). Its equation of state and its interaction with dark matter are in fact so far subject only to very weak constraints (Perlmutter et al. 1999, Huey et al. 1999, Baccigalupi et al. 2002, Corasaniti & Copeland 2002, Bean & Melchiorri 2002, Amendola et al. 2002) so that there exist a large variety of different dark energy models still viable (see for instance the review by Peebles & Ratra 2002). Perhaps the only characteristic common to almost all proposed models of dark energy is that its domination started very recently: the present epoch of acceleration was preceded by a decelerated epoch at redshift $z \geq 1$

in which structure formed. Two exceptions to this scheme are the model of Dodelson et al. 2000, in which the dark energy is periodically dominating, and the generic quintessence model of Lee & Ng 2002, in which the dark energy was dominating again at very large redshifts; in both cases the expansion was however decelerated at z of order unity. The fact that the acceleration sets in just recently in the cosmic history is one aspect of the “coincidence problem” that demands an explanation: see for instance the discussion based on the anthropic principle by Vilenkin (2001). Another aspect of the “coincidence problem” is commonly phrased as why the dark energy and the dark matter densities happen to be similar just today (Zlatev et al. 1999).

The existence of a decelerated epoch rests on three arguments, one theoretical and two observational. The theoretical argument is based on the most common models of dark energy. In fact, a FRW universe with scale factor $a = (1 + z)^{-1}$ filled with pressureless dark matter (subscript m) and non-interacting dark energy (subscript ϕ) with a constant equation of state $w_\phi = 1 + p_\phi/\rho_\phi$ is decelerated before the epoch

$$z_{acc} = [(2 - 3w_\phi)(\Omega_\phi/\Omega_m)]^{1/(3-3w_\phi)} - 1. \quad (1)$$

Here and in the following the density parameters Ω_i refer to the present quantity of the i -th component. Given the current estimates $\Omega_m \approx 0.3 \pm 0.1$, $\Omega_\phi = 0.7 \pm 0.1$ one has $z_{acc} \leq 1$ for all values of w . Even allowing for a large curvature, e.g. $|\Omega_k| \equiv |1 - (\Omega_m + \Omega_\phi)| < 0.3$, more than three sigma away from CMB measurements (de Bernardis et al. 2001), it is not possible to go beyond $z_{acc} \approx 1.3$. Therefore, an hypothetical observation of a value of z_{acc} significantly larger than unity would signal that some of the assumptions leading to (1) are false.

The two observational arguments are as follows. First, since gravitational instability is ineffective in an accelerated regime, it seems that an extended accelerated era is in contrast with the observed large scale structure. Second, the recent supernova SN1997ff at $z \approx 1.7$ is consistent with a decelerated expansion at the epoch of light emission (Benitez et al. 2001, Riess et al. 2002, Turner & Riess 2001) and seems to provide “a glimpse of the epoch of deceleration” (Benitez et al. 2001).

The aim of this paper is to show that all three arguments are not generally true: they are in fact valid only in a restricted class of models. As a counterexample, a simple flat-space model with a constant equation of state will be explicitly constructed that can be accelerated at large z , allows structure formation and is not in conflict with the current supernova data, including SN1997ff. Such a dark energy model, based on a coupling of dark energy to dark matter (Amendola 2000), makes very strong and unique predictions that can be easily tested in the near future.

Some of the results here reported have been already applied to a specific model of string-inspired dark energy (Amendola et al. 2002). This paper generalizes the arguments without referring to specific realizations and performs a likelihood comparison to the SNIa data.

2 THE $Z \approx 1.7$ SUPERNOVA

Let us start with the last argument, i.e. the constraints imposed by the farthest type Ia supernova known so far, SN1997ff (Benitez et al. 2001). The recent assessment of the lensing magnification increased the apparent magnitude of SN1997ff by 0.34 ± 0.12 mag (Riess et al. 2002). The result is conveniently expressed in terms of “Milne’s deviation”, i.e. the deviation $\Delta(m - M)$ of the distance modulus $m - M$ from that of a Milne model (i.e. a hyperbolic empty universe, $\Omega_m = \Omega_\Lambda = 0$ which has a constant expansion velocity). In Benitez et al. (2001) and Riess et al. (2002) it was found $\Delta(m - M) \approx -0.15 \pm 0.34$ at $z \approx 1.755$, in good agreement with a best fit flat $\Omega_m = 0.35, \Omega_\Lambda = 0.65$ cosmology. As we anticipated, for such a cosmology the redshift $z = 1.755$ is indeed already well within the decelerated epoch, which started at $z_{acc} = 0.548$. However, this does not mean that all models which are accelerated at large z are ruled out (even without mentioning the still unclear experimental uncertainties of the supernova detection). In fact, let us calculate the luminosity distance in Milne’s model. The luminosity distance in a FRW metric with scale factor $a = (1 + z)^{-1}$ is defined as

$$d_L(z) = \frac{1+z}{H_0} S \left[\int_0^z \frac{dz'}{E(z')} \right], \quad (2)$$

where the function $S[x]$ is $|\Omega_k|^{-1/2} \sin(|\Omega_k|^{1/2}x)$, x , $|\Omega_k|^{-1/2} \sinh(|\Omega_k|^{1/2}x)$ depending on the spatial curvature $k = 1, 0, -1$, respectively, and where the function $E(z)$ is derived from the Friedmann equation written as

$$H(z) = H_0 E(z). \quad (3)$$

For instance, in the case of a model with matter (subscript m), dark energy (subscript ϕ) with constant equation of state w_ϕ and curvature, one has

$$E^2(z) = \Omega_m(1+z)^3 + \Omega_\phi(1+z)^{3w_\phi} + \Omega_k(1+z)^2, \quad (4)$$

where $\Omega_k = 1 - \Omega_m - \Omega_\phi$. In an empty hyperbolic universe we have then $\Omega_k = 1$ and

$$E(z) = (1+z), \quad (5)$$

so that the luminosity distance is

$$d_{L(Milne)} = \frac{(1+z)}{H_0} S \left[\int_0^z \frac{dz'}{E(z')} \right] = \frac{z^2 + 2z}{2H_0}. \quad (6)$$

It appears then that $d_{L(Milne)} \sim z^2$ for large redshifts. Any model with a luminosity distance that grows slower than z^2 yields $\Delta(m - M) < 0$ (as SN1997ff) beyond some z .

Perhaps the simplest model one can construct with such a property is a flat space with a single fluid with constant equation of state w_e . In this case one has in fact

$$H^2 = H_0^2 a^{-3w_e}, \quad (7)$$




Figure 1. The thin curves represent the quantity $\Delta(m - M)$ for two values of w_e . The thick curves include a component $\Omega_b = 0.05$ of uncoupled baryons. The dashed curve is the reference model. The gray errorbars represents the high-redshift data of Perlmutter et al. (1999), while the black errorbars summarize all the supernova data at large z derived from Benitez et al. (2001).

and the luminosity distance (for $w \neq 2/3$)

$$d_L = (1 + z) \int_0^z \frac{dz'}{H(z')} = \frac{2(1 + z)}{(2 - 3w_e)H_0} \left[(1 + z)^{-\frac{3w_e}{2} + 1} - 1 \right]. \quad (8)$$

For $w_e < 2/3$ and large z we have $d_L \sim z^{2-3w_e/2}$. It follows that, at large z and for any $w_e > 0$, Milne's model gives always larger distances (and therefore fainter luminosity, i.e. larger apparent magnitudes) than the single-fluid flat-space evolution. In Fig. 1 I plot $\Delta(m - M)$ for various values of w_e , along with the reference flat model $\Omega_m = 0.35, \Omega_\Lambda = 0.65$. It appears clearly that values of w_e around 0.4 ± 0.1 are acceptable, as it will be confirmed by the likelihood analysis below.

However, according to the observations, our universe is composed by a mixture of, say, 30% of clustered matter (which includes a 5% of baryons) and 70% unclustered dark energy. Therefore the single-fluid model (7) can account for the observations only if both components scale with the same power w_e . This is exactly what happens assuming a direct interaction between dark matter and dark energy in the regime of strong coupling (Amendola 2000, Amendola & Tocchini-Valentini 2001, Gasperini et al. 2001). Let us assume infact two fluids: one (subscript m), with equation of state w_m , is later to be identified with dark matter, and the other (subscript ϕ), with

equation of state w_ϕ , to be identified with dark energy. Assume then a direct interaction term δ that transfers energy from one component to the other. The conservation equations are

$$\dot{\rho}_\phi + 3Hw_\phi\rho_\phi = -\delta, \quad (9)$$

$$\dot{\rho}_m + 3Hw_m\rho_m = \delta, \quad (10)$$

where the Friedmann equation reads

$$3H^2 = \kappa^2 (\rho_\phi + \rho_m), \quad (11)$$

and $\kappa^2 = 8\pi G$. It appears then that a solution with $\rho_m \sim \rho_\phi$ exists if (Zimdahl et al. 2001, Amendola & Tocchini-Valentini 2002)

$$\delta = 3H\Omega_\phi (w_m - w_\phi) \rho_m. \quad (12)$$

Therefore, the direct interaction induces on both components an effective equation of state

$$w_e = w_m + \Omega_\phi (w_\phi - w_m), \quad (13)$$

such that

$$\rho_m \sim \rho_\phi \sim a^{-3w_e}. \quad (14)$$

The Friedmann equation becomes then $H^2 = H_0^2 a^{-3w_e}$ as assumed in (7).

Accelerated cosmological solutions with $\rho_m \sim \rho_\phi$, denoted *stationary* solutions because $d(\rho_m/\rho_\phi)/dt = 0$, has been studied in several papers (see e.g. Amendola 1999, Chimento et al. 2000, Amendola & Tocchini-Valentini 2001, Gasperini et al. 2001, Pietroni 2002, Bonanno & Reuter 2002, Gromov et al. 2002) even before the evidences for acceleration (Carvalho et al. 1992, Wetterich 1995). This class of dark energy models has been invoked to explain the coincidence problem: in fact, in such models the dark matter and the dark energy scale identically with time, and as a consequence their ratio is frozen to the observed value from some time onward. It is indeed not difficult to produce accelerated expansions in which the ratio ρ_m/ρ_ϕ is of order unity at all times after equivalence (Amendola & Tocchini-Valentini 2001).

In Fig. 2 we plot the contours of the likelihood function obtained analysing the SNIa data of Perlmutter et al. (1999) to determine the allowed range of w_e (we consider the 38 high- z SNe plus the 16 low- z SNe of their fit C, plus the supernova at $z = 1.7$). In the upper part of the plot we show the likelihood $L(\Omega_m, w_\phi)$ for a “standard” flat-space model composed of two uncoupled fluid as in Eq. (4) with $\Omega_k = 0$: a fraction Ω_m of matter with equation of state $w_m = 1$ and a fraction $\Omega_\phi = 1 - \Omega_m$ of dark energy with constant equation of state w_ϕ . This plot reproduces the similar one in Perlmutter et al. (1999) (see also Dalal et al. 2001 for a further generalization) and has been obtained by marginalizing over the “nuisance” parameters M (the supernovae absolute magnitude) and α (the slope of the stretch factor relation, see Perlmutter et al. 1999 for the definitions of these parameters). In the limit of $\Omega_m \rightarrow 0$ the luminosity distance becomes as in the stationary case (8) with $w_\phi = w_e$. Therefore, in




Figure 2. Upper panel: marginalized likelihood function for a flat space model with two fluids, one with equation of state $w_m = 1$ and the other $w_\phi = \text{const.}$ The contours enclose 67%, 90% and 95%, dark to light gray. Lower panel: likelihood functions for the equation of state for a stationary model. Thick line: without baryons; thin line: including a fraction $\Omega_b = 0.05$ of uncoupled baryons; dashed line: including baryons and SN1997ff. The two horizontal lines mark the levels corresponding to 67% and 95% c.l.

the lower panel we show the one-dimensional likelihood $L(\Omega_m = 0, w_\phi)$, corresponding to the section at $\Omega_m = 0$ of the upper panel. As we will show in the following, it is necessary to decouple the baryons from the dark energy, so that the simple stationary case (8) has to be modified with the addition of a fraction Ω_b of uncoupled baryons. The likelihood $L(\Omega_b = 0.05, w_\phi)$ of this case is shown as a thin line. Finally, the intermediate likelihood drawn in dashed line includes also the supernova SN1997ff. The conclusion is that

$$w_e \in (0.2, 0.6) \quad (15)$$

at 95% c.l. ($w_e = 0.4 \pm 0.1$ at one sigma), with minimal variations including baryons and including SN1997ff. Notice that this is a one-parameter fit to the SNIa, with a $\chi^2/\text{d.o.f.} = 1.001$ for the best fit, just as good as the reference pure- Λ model.

Naturally, the stationary solution is acceptable only if one can show that the component ρ_m clusters while the dark energy component ρ_ϕ does not. This is the argument of the next section.

3 GROWTH OF PERTURBATIONS

Let us now identify the dark energy component with a scalar field. It has been shown in Amendola & Tocchini-Valentini (2001) that, assuming w_ϕ and w_m constant, the only scalar field model that reduces to the equations above requires a coupling

$$\delta = \sqrt{2/3\kappa\beta}|\dot{\phi}|\rho_m, \quad (16)$$

with

$$\beta = -\frac{3(w_m - 1 + \Omega_\phi(w_\phi - w_m))}{\sqrt{2(w_m - 1 + \Omega_\phi(1 - w_m + w_\phi))}} = \frac{3\Omega_\phi}{\sqrt{2w_\phi\Omega_\phi}}(1 - w_\phi), \quad (17)$$

where the second equality holds for pressureless dark matter, $w_m = 1$, which is the case we study below. Moreover, it is easy to show that $w_\phi = \text{const.}$ implies an exponential potential $U = U_0 e^{-\sqrt{2/3}\mu\kappa\phi}$ where

$$\mu = \frac{3(w_m + \Omega_\phi(w_\phi - w_m))}{\sqrt{2(w_m - 1 + \Omega_\phi(1 - w_m + w_\phi))}} = \frac{3}{\sqrt{2w_\phi\Omega_\phi}}[1 + \Omega_\phi(w_\phi - 1)] \quad (18)$$

(again the second equality is for $w_m = 1$). The equations for the matter (from now on $w_m = 1$) and the scalar field component are then

$$\ddot{\phi} + 3H\dot{\phi} + U_{,\phi} = -\sqrt{2/3\kappa\beta}\rho_m, \quad (19)$$

$$\dot{\rho}_m + 3H\rho_m = \sqrt{2/3\kappa\beta}\rho_m\dot{\phi}. \quad (20)$$

For $\beta > \sqrt{3}/2$ and $\mu > (-\beta + \sqrt{18 + \beta^2})/2$ (Amendola 2000), the system converges toward a global attractor with a power law expansion $a(t) \sim t^p$, characterized by

$$\Omega_\phi = \frac{4\beta^2 + 4\beta\mu + 18}{4(\beta + \mu)^2}, \quad (21)$$

$$w_e = \frac{\mu}{\mu + \beta}, \quad (22)$$

$$p = \frac{2}{3} \left(1 + \frac{\beta}{\mu} \right), \quad (23)$$

which is accelerated for $\mu < 2\beta$.

The growth of perturbations in the above model has been studied in Amendola & Tocchini-Valentini (2002); here we just state the results. For scales smaller than the horizon the perturbation equation for the matter density contrast δ_m can be written as

$$\delta_m'' + \left(2 + \frac{H'}{H} + \frac{3\beta}{\beta + \mu}\right) \delta_m' - \frac{3}{2}\gamma\delta_m\Omega_m = 0, \quad (24)$$

where primes denote derivation with respect to $\log a$ and $\gamma \equiv 1 + 4\beta^2/3$. Employing the trace of the Einstein equations

$$\frac{H'}{H} = -\frac{1}{2} \left[3 + \kappa^2 \left(\frac{1}{2}\phi'^2 - \frac{U}{H^2} \right) \right] = -\frac{3}{2}w_e, \quad (25)$$

we find the solution $\delta_m = a^{m_\pm}$ where

$$m_\pm = \frac{1}{4} \left[-4 + 3w_e - \frac{6\beta}{\beta + \mu} \pm \Delta \right], \quad (26)$$

where $\Delta^2 = 24\gamma\Omega_m + (-4 + 3w_e - \frac{6\beta}{\beta + \mu})^2$.

The observational constraints $\Omega_\phi = 0.7 \pm 0.1$ and $w_e = 0.4 \pm 0.1$ confine the parameters β, μ in the shadowed region in Fig. 3, where we also plot $m_+(\beta, \mu)$. For instance, putting $w_e = 0.4$, as suggested by the SNIa, and $\Omega_m = 0.3$ we obtain a growth as fast as $m_+ \approx 2$, which reduces to $m_+ \approx 1$ if $w_e = 0.5$. This shows that the dark matter fluctuations can grow even during the accelerated phase, due to the extra pull of the dark energy coupling.

The k -th plane-wave component of the scalar field perturbation $\varphi = \kappa(\delta\phi)/\sqrt{6}$ grows as

$$\varphi = \varphi_0 \delta_c \Omega_c (H_0/k)^2 a^{-3w_e+2}, \quad (27)$$

where

$$\varphi_0 = \beta - \frac{9}{\mu - 2\beta + (\beta + \mu)\Delta}. \quad (28)$$

For subhorizon wavelengths φ (which is proportional to $\delta\rho_\phi/\rho_\phi$) remains always much smaller than δ_m . It is interesting that even the dark energy fluctuations grow (as a^{m+2-3w_e}) during the accelerated epoch.

This demonstrates that the two components may be indeed identified with clustered matter and unclustered dark energy. Naturally, we have only proved that the perturbations do grow during acceleration, not that the present level of dark matter fluctuations is compatible with observations. This requires however the knowledge of the behavior of the perturbation throughout the post-decoupling era, not only during the final accelerated stage. In detailed models such as the string-inspired one discussed in Amendola et al. (2002) it can be shown that it is indeed possible to match the observed level of fluctuations.

As an aside, let us comments on the consequences of the above results on an application of the anthropic principle. According to Vilenkin (2002), that generalized previous analyses by Weinberg (1987), the amount of dark energy cannot exceed too much the observed value because otherwise the perturbation growth stops before galaxies can form, thereby preventing life. In particular, the anthropic principle




Figure 3. The shadowed region represents the observational constraints on β, μ . Within the unbroken curves $\Omega_\phi \in (0.6, 0.8)$; within the short-dashed curves $w_e \in (0.3, 0.5)$. The long-dashed curves represent $m(\beta, \mu) = 1$ (left curve) and $m(\beta, \mu) = 2$ (right curve).

“predicts” that Ω_ϕ has to be significantly smaller than unity, i.e. $\rho_\phi \approx \rho_m$. However, if dark energy is coupled to dark matter, perturbations can grow even when Ω_ϕ is dominating. For instance, a universe with $\Omega_\phi > 0.999$ (i.e. $\rho_\phi > 1000\rho_m$) admits a growth rate $m = 1$ if β is very large ($\beta > 50$ in this case). It seems therefore that the anthropic principle is unable to constrain Ω_ϕ in the general case.

4 WHEN DID THE ACCELERATION START?

We have shown so far that the past duration of the present acceleration is not limited by the supernovae data nor by the structure growth. But how far in the past can we extend the accelerated stationary solution? In this section we show that the limiting factor is to be found in the present observed abundance of baryons.

In fact, the coupling between matter and dark energy is a “fifth force” comparable in strength to gravity. This can be seen already from Eq. (24), in which the usual force term $3\delta_m\Omega_m/2 = 4\pi G(\delta\rho)_m/\rho_{crit}$ is replaced by $3\gamma\delta_m\Omega_m/2 = 4\pi G\gamma(\delta\rho)_m/\rho_{crit}$. In other words, to the lowest order, the effect of the dark matter/dark energy interaction is to induce an effective gravitational constant (Damour & Nordvedt 1993)

$$\tilde{G} = G\gamma = G \left(1 + \frac{4\beta^2}{3}\right). \quad (29)$$

As long as the dark energy interacts only with dark matter, such a fifth force is of course locally unobservable. However, if the dark energy couples to baryons with strength β_b then the interaction shows up to the post-post Newtonian (PPN) order as a non-zero parameter $\bar{\gamma}$ defined as in Hagiwara et al. (2002). The relation between $\bar{\gamma}$ and β_b can be found by observing that the coupled system of dark energy and matter fields ψ can be derived from the Lagrangian

$$L = \sqrt{|g|} \left(-\frac{R}{2\kappa^2} + \frac{1}{2} \phi_{;\mu} \phi^{;\mu} - V(\phi) \right) + L_m(\psi, \tilde{g}_{\mu\nu}), \quad (30)$$

where the metric $\tilde{g}_{\mu\nu} = e^{2a(\phi)} g_{\mu\nu}$ is conformally related to $g_{\mu\nu}$ by the factor

$$e^{2a(\phi)} = e^{-2\kappa\sqrt{\frac{2}{3}}\beta_b\phi} = e^{-4\sqrt{\frac{2}{3}}\beta_b\hat{\phi}}, \quad (31)$$

and where the dimensionless field $\hat{\phi} = \kappa\phi/2$ has been introduced to match the definition of Damour & Esposito-Farese (1992), Hagiwara et al. (2002) and Will (2002). Then, according to Damour & Esposito-Farese (1992) (see also Will 2002), the PPN parameter $\bar{\gamma}$ is defined as

$$\bar{\gamma} = -\frac{2\alpha_0^2}{1 + \alpha_0^2} = -\frac{4\beta_b^2}{3 + 2\beta_b^2}, \quad (32)$$

where $\alpha_0 = da(\phi)/d\phi$. The upper limits on $|\gamma|$ are of the order of 10^{-4} (Hagiwara et al. 2002), so that β_b is constrained to be smaller than 10^{-2} roughly. For as concerns the cosmological evolution at small z , this is equivalent to assuming $\beta_b = 0$, so that our starting assumption that the dark energy couples only (or preferentially) to dark matter, is justified. Similar species-dependent couplings have been discussed in other contexts since the first proposal by Damour, Gibbons, & Gundlach (1990), see e.g. the astrophysical bounds discussed by Gradwohl & Frieman (1992).

However, if the baryons are uncoupled they dilute with the usual behavior $\rho_b \sim a^{-3}$ i.e. faster than the coupled dark energy/dark matter fluid. This modifies the Friedmann equation (8) as follows

$$H^2 = H_0^2 [(1 - \Omega_b) a^{-3w_e} + \Omega_b a^{-3}]. \quad (33)$$

There appears therefore an epoch in the past before which the baryons were dominating and, as a consequence, the expansion decelerated. Denoting with Ω_u a generic uncoupled component (in this case the baryons) this epoch z_{acc} in flat space is

$$z_{acc}(\Omega_u, w_e) = [(2 - 3w_e)(1 - \Omega_u)/\Omega_u]^{1/(3-3w_e)} - 1, \quad (34)$$

which therefore replaces (1). It appears then that the factor that limits the acceleration epoch is the present abundance of baryons. Assuming $\Omega_u = \Omega_b \approx 0.05 \pm 0.02$ the maximum z_{acc} turns out to be around 5 as shown in Fig. 4.

It is to be noticed that if β and μ are constant at all epochs, the present accelerated




Figure 4. Contour plots of the function $z_{acc}(\Omega_u, w_e)$ where the uncoupled component is either the total matter content Ω_m , as in the standard dark energy models, or the baryon fraction Ω_b as in the model discussed in this paper. The observed total matter fraction is within the dashed lines, while the baryon fraction is within the dotted lines.

epoch is preceded by a decelerated baryon-dominated epoch before z_{acc} . Such an epoch would however be in conflict with the CMB, as shown in Tocchini-Valentini & Amendola (2001), because of an extremely large integrated Sachs-Wolfe effect on the CMB. It is therefore necessary to modify the simplest case with, for instance, a modulation of the coupling parameter β or the potential U in order to prevent the baryon domination, as in Amendola & Tocchini-Valentini (2001) and in Amendola et al. (2002). These models also account for the observed level of present fluctuations.

5 CONCLUSION

This paper shows that high- z acceleration is a viable possibility if dark energy couples to dark matter. Although the present abundance of baryons limit the epoch of acceleration to $z_{acc} < 5$, this is still much earlier than the standard models of uncoupled dark energy, which hardly reaches $z_{acc} = 1$. The future observations of SNIa at high redshift will be well suited to detect or reject an early acceleration.

A strong coupling between dark energy and dark matter can also be detected through the biasing and the rate of growth of perturbations, as discussed in Amendola & Tocchini-Valentini (2002). In principle, the coupling could be seen also in astrophysical objects like galaxy clusters by comparing the forces felt by dark matter halos and baryons (for instance, baryons in the intracluster gas), i.e. as an astrophysical test of the violation of the equivalence principle. This approach, discussed in detail in Gradwohl & Frieman (1992), requires several assumptions on the distribution of dark haloes. As emphasized by Peebles (2002), however, it has the potentiality to open unexplored paths in cosmology.

6 REFERENCES

- Amendola L. Phys. Rev. **D62**, 043511 (2000)
 Amendola L. and D. Tocchini-Valentini, Phys. Rev. **D64**, 043509 (2001)
 Amendola L. & D. Tocchini-Valentini, astro-ph/0111535, Phys. Rev. **D66**, 043528 (2002)
 Amendola L., C. Quercellini, D. Tocchini-Valentini and A. Pasqui, (2002) astro-ph/0205097
 Amendola L., M. Gasperini, D. Tocchini-Valentini and C. Ungarelli, (2002) astro-ph/0208032
 Baccigalupi C., A. Balbi, S. Matarrese, F. Perrotta, N. Vittorio, astro-ph/0109097, Phys.Rev. **D65** (2002) 063520
 Bahcall N. et al. 2002, astro-ph/0205490
 Bean R. & Melchiorri A., astro-ph/0110472, Phys.Rev. **D65** (2002) 041302
 Benitez N. et al., (2002) astro-ph/0207097
 Bonanno A. & M. Reuter (2002) Phys. Lett. B 527, 9
 Caldwell R.R., Dave R. & Steinhardt P.J. (1998), Phys. Rev. Lett. 80, 1582
 Carvalho J.C., J.A.S. Lima, I. Waga, Phys.Rev.**D46**:2404-2407,1992
 Chimento L.P., A. S. Jakubi & D. Pavon, Phys. Rev. **D62**, 063508 (2000), astro-ph/0005070;
 Corasaniti P.S.& E. Copeland, Phys.Rev. **D65** (2002) 043004
 Damour T.& Esposito-Farese G., Class. Quantum Grav. 9, 2093 (1992)
 Damour T. & Nordtvedt K., Phys. Rev. Lett. 70, 2217 (1993)
 Damour T., G. W. Gibbons and C. Gundlach, Phys. Rev. Lett., 64, 123, (1990)
 Dalal N. et al., Phys. Rev. Lett., **87**, 141302 (2001)
 De Bernardis et al. Nature 404:955-959,2000
 Dodelson S., M. Kaplinghat & E. Stewart, Phys. Rev. Lett. 85, 5276 (2000)
 Frieman J., C. T. Hill, A. Stebbins and I. Waga, Phys. Rev. Lett. **75**, 2077 (1995).
 Gasperini M., F. Piazza and G. Veneziano, Phys. Rev. **D65**, 023508 (2001).
 Gradwohl B.A.& Frieman, J. A. 1992 ApJ, 398, 407
 Gromov A., Yu. Baryshev & P. Teerikorpi (2002) astro-ph/0209458
 Hagiwara K. et al., Phys. Rev. **D66** 010001-1 (2002), available at pdg.lbl.gov
 Halverson N.W.*et al.*, (2001) astro-ph/0104489.

- Huey G., Wang, L., Dave, R., Caldwell, R. R., & Steinhardt, P. J. (1999), Phys. Rev. D **59**, 063005.
- Lee A.T.*et al.*, Ap. J. **561**, L1 (2001).
- Lee W.& K.-W. Ng astro-ph/0209093.
- Netterfield C.B.*et al.*, astro-ph/0104460
- Perlmutter S.*et al.*, Ap. J. **517**, 565 (1999).
- Peebles P.J.E.& B. Ratra, astro-ph/0207347
- Peebles P.J.E., astro-ph/0208037
- Pietroni M., hep-ph/0203085.
- Ratra B.& P.J.E. Peebles, Phys. Rev. D **37**, 3406 (1988)
- Riess A.*et al.*, Astron. J. **116**, 1009 (1998).
- Riess A. et al. Ap. J. **560**, 49 (2001)
- Tocchini-Valentini D. and L. Amendola, Phys.Rev. **D65**, 063508 (2002)
- Turner M. and A. Riess, astro-ph/0106051
- Vilenkin A., hep-th/0106083
- Weinberg S. Phys. Rev. Lett. **59**, 2607 (1987)
- Wetterich C. (1988) Nucl. Phys. B., **302**, 668;
- Wetterich C., Astronomy and Astrophys. **301**, 321 (1995).
- Will C., Living Rev. Relativity, **4** (2001), 4: cited on Sep. 19, 2002, <http://www.livingreview.org/Articles/V4will/>
- Zimdahl W., D. Pavon & L. P. Chimento, Phys.Lett. B **521** (2001) 133, astro-ph/0105479
- Zlatev I., L. Wang & P.J. Steinhardt, Phys. Rev. Lett. **82** 896 (1999).